

NEWS 25 APR 30 CHEMCATS enhanced with 1.2 million new records  
 NEWS 26 APR 30 CA/CAPplus enhanced with 1870-1889 U.S. patent  
 records  
 NEWS 27 APR 30 INPADOC replaced by INPADOCDB on STN  
 NEWS 28 MAY 01 New CAS web site launched  
 NEWS 29 MAY 08 CA/CAPplus Indian patent publication number format  
 defined  
 NEWS 30 MAY 14 RDISCLOSURE on STN Easy enhanced with new search  
 and display fields  
 NEWS 31 MAY 21 BIOSIS reloaded and enhanced with archival data  
 NEWS 32 MAY 21 TOXCENTER enhanced with BIOSIS reload  
 NEWS 33 MAY 21 CA/CAPplus enhanced with additional kind codes for  
 German patents  
 NEWS 34 MAY 22 CA/CAPplus enhanced with IPC reclassification in  
 Japanese patents  
 NEWS EXPRESS NOVEMBER 10 CURRENT WINDOWS VERSION IS V8.01c, CURRENT  
 MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP),  
 AND CURRENT DISCOVER FILE IS DATED 25 SEPTEMBER 2006.  
 NEWS HOURS STN Operating Hours Plus Help Desk Availability  
 NEWS LOGIN Welcome Banner and News Items  
 NEWS IPC8 For general information regarding STN implementation  
 of IPC 8

Enter NEWS followed by the item number or name to see news on that  
 specific topic.

All use of STN is subject to the provisions of the STN Customer  
 agreement. Please note that this agreement limits use to scientific  
 research. Use for software development or design or implementation  
 of commercial gateways or other similar uses is prohibited and may  
 result in loss of user privileges and other penalties.

\* \* \* \* \* STN Columbus \* \* \* \* \*  
 \* \*

FILE 'HOME' ENTERED AT 16:24:14 ON 11 JUN 2007

=> File Medline EMBASE Biosis Caplus  
 COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
0.21	0.21

FULL ESTIMATED COST

FILE 'MEDLINE' ENTERED AT 16:24:58 ON 11 JUN 2007

FILE 'EMBASE' ENTERED AT 16:24:58 ON 11 JUN 2007

Copyright (c) 2007 Elsevier B.V. All rights reserved.

FILE 'BIOSIS' ENTERED AT 16:24:58 ON 11 JUN 2007  
Copyright (c) 2007 The Thomson Corporation

FILE 'CAPLUS' ENTERED AT 16:24:58 ON 11 JUN 2007  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

=> s (casein kinase) (w) (I or 1)  
L1 2877 (CASEIN KINASE) (W) (I OR 1)

=> s (inhibitor or inhibitors)  
L2 2755119 (INHIBITOR OR INHIBITORS)

=> s L1 (4A) L2  
L3 136 L1 (4A) L2

=> s L2 (6A) cell or cellular  
L4 1591079 L2 (6A) CELL OR CELLULAR

=> s L3 and L4  
L5 17 L3 AND L4

=> duplicate  
ENTER REMOVE, IDENTIFY, ONLY, OR (?):remove  
ENTER L# LIST OR (END):15  
DUPLICATE PREFERENCE IS 'MEDLINE, EMBASE, BIOSIS, CAPLUS'  
KEEP DUPLICATES FROM MORE THAN ONE FILE? Y/(N):n  
PROCESSING COMPLETED FOR L5  
L6 10 DUPLICATE REMOVE L5 (7 DUPLICATES REMOVED)

=> d L6 1-10 bib ab

L6 ANSWER 1 OF 10 CAPLUS COPYRIGHT 2007 ACS on STN  
AN 2006:13464 CAPLUS  
DN 144:101073  
TI therapeutic uses of kinase inhibitors, and compositions thereof  
IN Caligiuri, Maureen G.; Kley, Nikolai A.; Murthi, Krishna K.  
PA GPC Biotech, Inc., USA  
SO PCT Int. Appl., 201 pp.  
CODEN: PIXXD2  
DT Patent  
LA English  
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.
DATE	-----	----	-----	-----
-----				
PI	WO 2006002119	A2	20060105	WO 2005-US21843
	20050617			

WO 2006002119 A3 20070222  
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ,  
CA, CH,  
CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,  
GB, GD,  
GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP,  
KR, KZ,  
LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,  
MZ, NA,  
NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE,  
SG, SK,  
SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC,  
VN, YU,  
ZA, ZM, ZW  
RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR,  
HU, IE,  
IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF,  
BJ, CF,  
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW,  
GH, GM,  
KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ,  
BY, KG,  
KZ, MD, RU, TJ, TM

EP 1763345 A2 20070321 EP 2005-762859  
20050617  
R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR,  
HU, IE,  
IS, IT, LI, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR,  
AL, BA,  
HR, LV, MK, YU

PRAI US 2004-580868P P 20040618  
WO 2005-US21843 W 20050617

OS MARPAT 144:101073

AB The invention pertains to inhibitors of various kinases (e.g. S/T kinases,

Tyr kinases, etc.), which inhibitors are previously known as cyclin

dependent kinase inhibitors (CDKs). The inhibitors of the invention are

capable of inhibiting various wild-type and mutant form kinases, including

drug-resistant forms of mutant kinases. Thus, the kinase inhibitors are

useful in treating a wide range of diseases/conditions associated with

abnormal functions/excessive activities of the target kinases, including

mutant kinases. The invention further provides methods for treating

cancers, tumors and patients which are resistant or refractory to other

therapeutic agents. Pharmaceutical compns. and packaged pharmaceuticals with instructions of these inhibitors, and methods of using these inhibitors are also provided.

L6 ANSWER 2 OF 10 MEDLINE on STN DUPLICATE 1  
AN 2006124588 MEDLINE  
DN PubMed ID: 16247451  
TI RNAi-based screening of the human kinome identifies  
Akt-cooperating  
kinases: a new approach to designing efficacious multitargeted  
kinase  
inhibitors.  
AU Morgan-Lappe S; Woods K W; Li Q; Anderson M G; Schurdak M E; Luo  
Y;  
Giranda V L; Fesik S W; Levenson J D  
CS Abbott Laboratories, Cancer Research, Abbott Park, IL 60064,  
USA.  
SO Oncogene, (2006 Mar 2) Vol. 25, No. 9, pp. 1340-8.  
Journal code: 8711562. ISSN: 0950-9232.  
CY England: United Kingdom  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 200604  
ED Entered STN: 3 Mar 2006  
Last Updated on STN: 19 Apr 2006  
Entered Medline: 18 Apr 2006  
AB Tumors comprise genetically heterogeneous cell populations,  
whose growth  
and survival depend on multiple signaling pathways. This has  
spurred the  
development of multitargeted therapies, including small  
molecules that can  
inhibit multiple kinases. A major challenge in designing such  
molecules  
is to determine which kinases to inhibit in each cancer to  
maximize  
efficacy and therapeutic index. We describe an approach to this  
problem  
implementing RNA interference technology. In order to identify  
Akt-cooperating kinases, we screened a library of  
kinase-directed small  
interfering RNAs (siRNAs) for enhanced cancer cell killing in  
the presence of Akt inhibitor A-443654. siRNAs targeting  
casein kinase I gamma 3 (CSNK1G3) or the  
inositol polyphosphate multikinase (IPMK) significantly enhanced  
A-443654-mediated cell killing, and caused decreases in Akt  
Ser-473 and  
ribosomal protein S6 phosphorylation. Small molecules targeting  
CSNK1G3

and/or IPMK in addition to Akt may thus exhibit increased efficacy and have the potential for improved therapeutic index.

L6 ANSWER 3 OF 10 CAPLUS COPYRIGHT 2007 ACS on STN  
AN 2005:451232 CAPLUS  
DN 143:19954  
TI Methods for inhibiting cell growth  
IN Zhao, Yi; Chandraratna, Roshantha A.  
PA Allergan, Inc., USA  
SO PCT Int. Appl., 78 pp.  
CODEN: PIXXD2  
DT Patent  
LA English  
FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.
DATE			
-----	----	-----	-----
PI WO 2005046726	A2	20050526	WO 2004-US37881
20041112			
WO 2005046726	A3	20051208	
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
PRAI US 2003-519528P	P	20031112	
US 2004-564807P	P	20040422	
AB Cell growth is inhibited and/or cell death is induced in a cell by administering an RXR (retinoid X receptor) agonist and an inhibitor of casein kinase 1			
α. A cell or a tissue can be screened for enhanced susceptibility			

to cell death or interference with cell growth. Conditions characterized by uncontrolled cell growth or proliferation, such as a cancer, can be treated with inhibitors of casein kinase 1  $\alpha$ .

L6 ANSWER 4 OF 10 CAPLUS COPYRIGHT 2007 ACS on STN  
 AN 2002:521462 CAPLUS  
 DN 137:88442  
 TI Incensole and furanogermacrene and compounds in treatment for inhibiting neoplastic lesions and microorganisms  
 IN Shanahan-Pendergast, Elisabeth  
 PA Ire.  
 SO PCT Int. Appl., 68 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.
WO 2002053138	A2	20020711	WO 2002-IE1
WO 2002053138	A3	20020919	
W: AE, AG, AT, AU, BB, BG, CA, CH, CN, CO, CU, CZ, LU, LV, MA, MD, UA, UG, US, VN, YU, RU, TJ, TM			
RW: GH, GM, KE, LS, MW, SD, SL, SZ, UG, AT, BE, CH, CY, DE, ES, FI, ML, MR, NE, SN, TD, TG			
AU 2002219472	A1	20020716	AU 2002-219472
EP 1351678	A2	20031015	EP 2002-727007
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			
US 2004092583	A1	20040513	US 2004-250535
IE 2001-2	A	20010102	
WO 2002-IE1	W	20020102	
AB The invention discloses the use of incensole and/or furanogermacrene, derivs. metabolites and precursors thereof in the treatment of neoplasia, particularly resistant neoplasia and immunodysregulatory disorders. These			

compds. can be administered alone or in combination with conventional  
chemotherapeutic, antiviral, antiparasite agents, radiation  
and/or surgery. Incensole and furanogermacren and their mixture showed  
antitumor activity against various human carcinomas and melanomas and  
antimicrobial activity against Staphylococcus aureus and Enterococcus faecalis.

L6 ANSWER 5 OF 10 CAPLUS COPYRIGHT 2007 ACS on STN  
AN 2001:432817 CAPLUS  
DN 135:41041  
TI Use of hymenialdisine or a derivative thereof as an inhibitor of  
cyclin-dependent kinases, GSK-3 $\beta$  and casein kinase 1, and  
therapeutic use  
IN Meijer, Laurent  
PA Centre National de la Recherche Scientifique (CNRS), Fr.  
SO Eur. Pat. Appl., 38 pp.  
CODEN: EPXXDW  
DT Patent  
LA English  
FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.
EP 1106180	A1	20010613	EP 1999-403077
19991208			
EP 1106180	B1	20031112	
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,			
IE, SI, LT, LV, FI, RO			
AT 253918	T	20031115	AT 1999-403077
19991208			
ES 2213996	T3	20040901	ES 1999-403077
19991208			
CA 2384982	A1	20010614	CA 2000-2384982
20001207			
WO 2001041768	A2	20010614	WO 2000-EP12791
20001207			
WO 2001041768	A3	20020510	
WO 2001041768	A9	20020912	
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,			
CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,			
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT,			
LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU,			

SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US,  
UZ, VN,

YU, ZA, ZW

RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE,  
CH, CY,

DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE,  
TR, BF,

BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG

EP 1235578 A2 20020904 EP 2000-987404  
20001207

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,  
MC, PT,

IE, SI, LT, LV, FI, RO, MK, CY, AL, TR

JP 2004500356 T 20040108 JP 2001-543113  
20001207

US 2003105075 A1 20030605 US 2002-149115  
20021004

US 7098204 B2 20060829

PRAI EP 1999-403077 A 19991208

WO 2000-EP12791 W 20001207

AB The title compds. are I (R1, R2 = H, Br), or a pharmaceutically  
acceptable

salt thereof, are used for the manufacture of a medicament for  
inhibiting

cyclin-dependent kinases, GSK-3 $\beta$ , and casein kinase 1. The  
compds.

may be used for preventing and treating neurodegenerative  
disorders (e.g.

Alzheimer's disease), diabetes, inflammatory pathologies, and  
cancers.

RE.CNT 13 THERE ARE 13 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 6 OF 10 CAPLUS COPYRIGHT 2007 ACS on STN

AN 2000:856756 CAPLUS

DN 134:129061

TI IC261, a specific inhibitor of the protein kinases  
casein kinase 1-delta and -epsilon, triggers  
the mitotic checkpoint and induces p53-dependent postmitotic  
effects

AU Behrend, L.; Milne, D. M.; Stoter, M.; Deppert, W.; Campbell, L.  
E.; Meek,

D. W.; Knippschild, U.

CS Heinrich-Pette-Institut fur Experimentelle Virologie und  
Immunologie,

Hamburg, D-20251, Germany

SO Oncogene (2000), 19(47), 5303-5313

CODEN: ONCNES; ISSN: 0950-9232

PB Nature Publishing Group

DT Journal

LA English



AB The p53-targeted kinases casein kinase 1δ (CK1δ) and casein kinase 1ε (CK1ε) have been proposed to be involved in regulating DNA repair and chromosomal segregation. Recently, we showed that CK1δ localizes to the spindle apparatus and the centrosomes in cells with mitotic failure caused by DNA-damage prior to mitotic entry. We provide here evidence that 3-(2,4,6-trimethoxyphenyl)methylidenyl-indolin-2-one (IC261), a novel inhibitor of CK1δ and CK1ε, triggers the mitotic checkpoint control. At low micromolar concns. IC261 inhibits cytokinesis causing a transient mitotic arrest. Cells containing active p53 arrest in the postmitotic G1 phase by blockage of entry into the S phase. Cells with non-functional p53 undergo postmitotic replication developing an 8N DNA content. The increase of DNA content is accompanied by a high amount of micronucleated and apoptotic cells. Immunofluorescence images show that at low concns. IC261 leads to centrosome amplification causing multipolar mitosis. Our data are consistent with a role for CK1δ and CK1ε isoforms in regulating key aspects of cell division, possibly through the regulation of centrosome or spindle function during mitosis.

RE.CNT 67 THERE ARE 67 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 7 OF 10 MEDLINE on STN DUPLICATE 2  
AN 1998366074 MEDLINE  
DN PubMed ID: 9700717  
TI H-7-induced apoptosis in the cells of a Drosophila neuronal cell line through affecting unidentified H-7-sensitive substance(s).  
AU Nagano M; Suzuki H; Ui-Tei K; Sato S; Miyake T; Miyata Y  
CS Department of Pharmacology, Nippon Medical School, Tokyo, Japan.  
SO Neuroscience research, (1998 Jun) Vol. 31, No. 2, pp. 113-21.  
Journal code: 8500749. ISSN: 0168-0102.  
CY Ireland  
DT (COMPARATIVE STUDY)  
Journal; Article; (JOURNAL ARTICLE)  
(RESEARCH SUPPORT, NON-U.S. GOV'T)  
LA English  
FS Priority Journals  
EM 199811

ED Entered STN: 6 Jan 1999  
 Last Updated on STN: 6 Jan 1999  
 Entered Medline: 20 Nov 1998

AB The present study was undertaken to reveal underlying mechanisms of apoptosis in neurons using clonal neuronal cells, ML-DmBG2-c2, derived from *Drosophila* larval central nervous system

1-(5-Isoquinolinesulfonyl)-2-methylpiperazine (H-7), a protein kinase inhibitor, induced cell death with typical features of apoptosis such as internucleosomal DNA fragmentation, nuclear condensation and apoptotic bodies in the cells. Though H-7 is known to inhibit cAMP-dependent protein kinase (PKA), protein kinase C (PKC), cGMP-dependent kinase (PKG), myosin light chain kinase (MLCK), and casein kinase I (CKI), specific inhibitors for these kinases such as H-89, calphostin C, ML-9, or CKI-7 did not induce apoptosis in the cells. Other kinases such as tyrosine kinase. PI3-kinase and Ca<sup>2+</sup>/CaM kinase II so far examined in the present study were interpreted not to be involved in the apoptotic cascade. Therefore, it is concluded that an H-7-sensitive substance(s) other than these kinases is responsible for the apoptosis in the neuronal cells.

Caspase inhibitors prevented apoptosis in the cells treated with H-7. These results suggest that caspase(s) is involved downstream of the H-7-sensitive point in the cascade of the apoptosis.

L6 ANSWER 8 OF 10 CAPLUS COPYRIGHT 2007 ACS on STN  
 AN 1995:301667 CAPLUS  
 DN 122:127030  
 TI Development of inhibitors of protein kinases CKI and CKII and some related aspects, including donor and acceptor specificities and viral protein kinases

AU Shugar, David  
 CS Inst. Biochem. Biophysics, Polish Academy Sciences, Warszawa, 02-106, Pol.  
 SO Cellular & Molecular Biology Research (1994), 40(5/6), 411-19  
 CODEN: CMBREW; ISSN: 0968-8773  
 PB Elsevier  
 DT Journal; General Review  
 LA English  
 AB A review with .apprx.45 refs. A brief overview is presented of progress

in the development of specific inhibitors of protein kinases CKI and CKII.

Two promising classes of inhibitors, which have the ability to traverse cell membranes, are now known. One of these is based on halogenated benzimidazoles and 2-aza-benzimidazoles (benzotriazoles)

and some of their nucleosides. The second embraces modified isoquinoline

sulfonamides, several of which are known as inhibitors of other protein

kinases. Both classes include analogs that permit discrimination between

CKI and CKII. Ongoing research with halogenated benzotriazoles leads to

inhibitors with  $K_i$  values below 1  $\mu$ M. Also considered are nucleoside

triphosphate analog inhibitors and their potential properties as donors,

with illustrative examples from the field of nucleoside kinases, including

the apparent existence of a dual-specific viral protein/nucleoside kinase.

The role of cellular CKII and viral-encoded CKII-like activities in viral replication underlines the potential of CKII inhibitors as

antiviral agents, exemplified by the case of vesicular stomatitis virus.

L6 ANSWER 9 OF 10 MEDLINE on STN

DUPLICATE 3

AN 91120135 MEDLINE

DN PubMed ID: 2278876

TI A protein complex expressed during terminal differentiation of monomyelocytic cells is an inhibitor of cell growth.

AU Murao S; Collart F; Huberman E

CS Biological and Medical Research Division, Argonne National Laboratory,  
Illinois 60439.

SO Cell growth & differentiation : the molecular biology journal of the

American Association for Cancer Research, (1990 Oct) Vol. 1, No. 10, pp.

447-54.

Journal code: 9100024. ISSN: 1044-9523.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)

(RESEARCH SUPPORT, U.S. GOV'T, NON-P.H.S.)

LA English

FS Priority Journals

EM 199103

ED Entered STN: 5 Apr 1991

Last Updated on STN: 3 Feb 1997

Entered Medline: 12 Mar 1991

AB A protein complex (PC) composed of the MRP8 and MRP14 proteins has

previously been shown to be a specific inhibitor of casein kinase I and II. This PC is expressed during the late stages of terminal differentiation induced in

human

promyelocytic HL-60 leukemia cells by 1

alpha,25-dihydroxyvitamin D3 and

in human monocytic THP-1 leukemia cells by phorbol 12-myristate 13-acetate. This expression is associated with terminal cell differentiation because incubation of HL-60 cells with an agent

or

condition that causes suppression of growth but not induction of differentiation does not result in expression of the PC. At concentrations of 5-15 nM, the purified PC inhibited the growth

of HL-60

cells and THP-1 cells, as well as other cell types belonging to different

cell lineages. This growth inhibition was preceded by a

reduction in

[32P]phosphate incorporation and, at the higher PC

concentrations, was

associated with a reduction in [3H]thymidine, [3H]uridine, and

[32S]methionine incorporation. The specific expression pattern

and

growth-inhibitory character of the PC suggests that the complex may have a

role in suppressing cell growth during monomyelocytic terminal differentiation induced by specific chemical stimuli and during physiological and pathological events associated with

monomyelocytic cell

functions.

L6 ANSWER 10 OF 10 CAPLUS COPYRIGHT 2007 ACS on STN

AN 1990:607124 CAPLUS

DN 113:207124

TI Casein kinase 2: an 'eminence grise' in cellular regulation?

AU Pinna, Lorenzo A.

CS Dip. Chim. Biol., Univ. Padova, Padua, 35121, Italy

SO Biochimica et Biophysica Acta, Molecular Cell Research (1990), 1054(3),

267-84

CODEN: BBAMCO; ISSN: 0167-4889

DT Journal; General Review

LA English

AB A review, with 176 refs., on casein kinase 2 (CK2) with emphasis on the

features of CK2, subunit composition, structure of the  $\alpha$ - and  $\beta$ -subunits, regulation of CK2, biol. functions, phosphorylatable substrates, substrate and inhibitor specificity, and comparison

to casein kinase 1.